

The following book chapter 20 might be cited as:

Dokulil MT, Donabaum K, Pall K, Janauer GA, Teubner K (2018) Eutrophication, Management and Sustainable Development of Urban Lakes: General Considerations and Specific Solutions for Alte Donau – A Synthesis. pp 373-385. *In*: Dokulil MT, Donabaum K, Teubner K (eds) The Alte Donau: Successful Restoration and Sustainable Management. Aquatic Ecology Series, vol 10. Springer
(doi: [10.1007/978-3-319-93270-5_20](https://doi.org/10.1007/978-3-319-93270-5_20))

The final publication is available at Springer via:

https://doi.org/10.1007/978-3-319-93270-5_20

Aquatic Ecology Series

Martin T. Dokulil · Karl Donabaum
Katrin Teubner *Editors*

The Alte Donau: Successful Restoration and Sustainable Management

An Ecosystem Case Study of a Shallow
Urban Lake

 Springer

Chapter 20

Eutrophication, Management and Sustainable Development of Urban Lakes: General Considerations and Specific Solutions for Alte Donau – A Synthesis

AUT Martin T. Dokulil, Karl Donabaum, Karin Pall, Georg A. Janauer, and Katrin Teubner

Abstract Intensively used urban water bodies are vulnerable to eutrophication. The shallow lake Alte Donau (Vienna) can be seen as an example for the extent of anthropogenic influence. Human impacts paired with changes in environmental conditions gave way to eutrophication processes in Alte Donau. Due to the great public interest restoration concepts and subsequently management programs were established. This chapter provides a synthesis of the key aspects to evolve and implement a successful water management plan. An attempt is made to generalise our specific solutions to serve as a basis for the development of similar strategies for other urban lakes.

Keywords Restoration · Groundwater · Seepage · Management · Improvement · Shallow lake

K. Donabaum · M. T. Dokulil (✉)
DWS Hydro-Ökologie, Wien, Austria
e-mail: karl.donabaum@dws-hydro-oekologie.at; martin.dokulil@univie.ac.at

K. Pall
Systema GmbH, Wien, Austria
e-mail: karin.pall@systema.at

K. Teubner · G. A. Janauer
Department of Limnology & Biological Oceanography, University of Vienna, Vienna, Austria
e-mail: katrin.teubner@univie.ac.at; georg.janauer@univie.ac.at

20.1 Introduction

Urban lakes are different types of natural or man-made water bodies within densely populated areas. For the purposes of management, these lakes can be defined by several operational criteria. They tend to be small and shallow, with surface areas of less than 2.5 km² and a mean depth of 6 m or less. The ratio of the watershed area to lake area can be variable but is often 10:1 or higher, meaning that their watersheds exert a strong influence on the lake. As an overall index of development, the urban lake watershed must contain at least 5% impermeable cover. The water budget often depends on ground-water and precipitation or is entirely artificially controlled. Most of these urban lakes must be managed for recreation, water supply, flood control or some other direct human use regardless whether natural or man-made (Birch and Mc Caskie 1999).

Curiously, the unique problems and conditions of urban lakes have received little attention in scientific literature. This is particularly surprising given that many of the management efforts are devoted to lakes and reservoirs that are distinctly urban in character. While the watershed management literature is filled with phosphorus budgets and watershed models, it is unusual to find overviews about the influence of watershed development on lake quality and it is exceptionally rare to find studies that have tracked changes in lake quality as a function of watershed development over time (Schueler and Simpson 2001).

However, restoration and recovery of already eutrophicated systems can only be achieved if external measures like watershed management are combined with internal measures.

20.2 Eutrophication

Small and shallow aquatic ecosystems generally have lower resilience than large, deep lakes. Urban lakes are therefore more sensitive to water pollution and eutrophication suffering from natural or anthropogenic impacts. If man-made, these environments tend to have rather regular shapes and higher shoreline development when recreationally used resulting in negative impacts on their functioning (Naselli-Flores 2008).

Metropolitan runoff flowing over impermeable surfaces collects large amounts of nutrients resulting in higher unit area phosphorus load from storm-water than other watersheds. Many urban watersheds receive additional loads from storm-water overflow, failing septic systems or pollutant seepage. Urban lakes also have unique internal nutrients sources such as water bird droppings, boat sewage and sediment release (Traut and Hostetler 2004). As a consequence, phytoplankton blooms and uncontrolled macrophyte growth may severely impair water quality and cause sanitary risks. Massive growth of submerged plants may form an obstacle to several forms of recreational use. Moreover, highly developed shorelines, including

tourism exploitation (Dokulil 2014a), housing and development sites (Cappiella and Schueler 2001) may significantly contribute to eutrophication and pollution. Urban lakes may potentially be contaminated by various compounds particularly their sediments causing long-term environmental problems and human health risks. Because of these many facets impacts have on urban lakes, management efforts must be integrative and sustainable (e.g. Sorensen 1996) especially when impacts are amplified by climate warming (Dokulil 2014b).

20.3 Restoration and Management

Urban lakes are complex systems strongly influenced by disturbances within their watershed. Therefore, thorough management of urban lakes must be tightly coupled with the management of the watershed. The entire catchment must be supervised and necessary strategies developed to optimise land use, erosion, housing, wastewater treatment, public transport, recreation, tourism or any other factor which might be important in the watershed (NALMS 1988).

Internal restoration techniques are abundantly described particularly for shallow water bodies (Carvalho 1994; Xu et al. 1999; Morscheid and Maehlmann 2005). Methods have been summarised and classified in Fig. 20.1 after Singh (1982). Among the many methods available, chemical stabilisation of phosphorus (Welch and Schrieve 1994) and removal of lake sediment (Van der Does and Frinking 1993; Björk 1978, 1994) are most popular. In many food web oriented lake rehabilitation




Fig. 20.1 Classification of lake restoration techniques. (Modified from Singh 1982)

activities, manipulation of the fish community is a prime focus (Hansson and Butler 1994; Berg et al. 1997; Hansel-Welch et al. 2003).

As aquatic plants may easily become a nuisance in man-made waters, management of aquatic vegetation is a major requisite usually by controlling abundant growth of macrophytes, e.g. by extensive harvesting (Pieterse and Murphy 1990; Bowmer et al. 1984; Madsen et al. 1988). More recently, macrophytes are recognised as an essential component in suppressing algal dominance during rehabilitation of eutrophic water bodies (Körner 2002). Regrowth of macrophytes was reported in a majority of lake restoration studies (Hansson and Butler 1994; Perrow et al. 1997). The re-establishment of aquatic vegetation in these studies occurred mainly by natural propagation and without any active management (Hansel-Welch et al. 2003). Under such conditions, the recovery of the water body towards the macrophyte dominated state may take long and the ecosystem is at risk to shift back to the algal dominated state.

Investigations on the essential environmental conditions promoting macrophyte growth are numerous (e.g. Riis and Hawes 2002; Crisman et al. 2005; Janse et al. 1998). The complexity of ecological pressures in lake environments, ranging from sediment characteristics to herbivory, often impairs the proliferation of the aquatic plants (Lau and Lane 2002; Irfanullah and Moss 2004).

The submersed aquatic vegetation is the essential factor determining water quality by preventing the growth of phytoplankton in shallow lakes (Bailey et al. 2002; Gopal and Goel 1993; Wium-Andersen 1987; Berger and Schagerl 2003).

Several experimental approaches were developed for the transplantation of submersed plants for pools (Irfanullah and Moss 2004), lake littorals (Wyckera and Humpesch 2002; IGB 2005; Hilt 2005; Gross and Hilt 2005; Morscheid and Maehlmann 2005) and for oxbows (Janauer 1995; Janauer and Pall 1998).

20.4 Sustainable Development

Water is so essential to life and the life processes of all living beings, that management of water resources requires a new paradigm, the concept of sustainability (Heintz 2004; Taylor and Goldstein 2010). This concept describes the dynamic conditions and the resiliency or robustness of complex systems to adapt and thrive in the face of change. Sustainable development shall meet the needs of the present without compromising vital ecosystems as well as the ability of future generations to meet their own needs. Application of the concept to water resources management involves integrative components. Once sustainable goals have been defined and adopted, sustainable infrastructure principles can operate. These principles must be based on the best integrated technology and institutional capacity. The outcome must then be evaluated and adapted if necessary in an iterative process. All activities in sustainable development of water resources shall rely on basin wide perspectives.




Fig. 20.2 Annual changes of macrophyte biomass (BM) in t dry mass (bars) and phytoplankton biovolume (B) as mg fresh weight per litre (line) from 1987 to 2011. Arrows with continuous line: Riplox-treatment, arrow with dashed line: start of periodical water level lowering

20.5 Specific Solutions for Alte Donau

118

Restoration and management measures for sustainable development of the urban lake Alte Donau were extensively described and discussed in this volume. The key measures and their results shall be briefly summarised here.

119
120
121

Excessive nutrient concentrations were strongly reduced by phosphorus flocculation with ferric chloride and by sediment oxidation with calcium nitrate (Chap. 5 – Riplox treatment, Chap. 6). The efficacy of this treatment was predicted to be 10 years. Algal blooms disappeared and across the years phytoplankton composition shifted from blue-greens to green algae, diatoms and later on to chrysophytes (Chaps. 9 and 10). Concomitantly the metazoan zooplankton altered from mainly filter-feeding herbivorous cladocerans under eutrophic algal-turbid state to mainly selective-feeding omnivorous and herbivorous copepods under mesotrophic transparent-water state (Chap. 11). Assemblages of microzoans (rotifers, Chap. 11) and protozoans (ciliates in Chap. 12, heterotrophic nanoflagellates in Chap. 13) differed between treatment periods by lowered food supply (see also bacteria in Chap. 13 and discussion Chap. 11) or increasing grazing pressure.

122
123
124
125
126
127
128
129
130
131
132
133

With the reduction of the phosphorus concentration in the water column and associated suppression of phytoplankton development, the transparency increased significantly (Chap. 6) promoting an initial recovery of macrophyte-stands (Fig. 20.2). But in the following years the macrophytes showed no significant increase in biomass, even though regrowth was assisted by plantings. Figure 20.3 shows that the hysteresis of the year by year trajectory indicates considerable

134
135
136
137
138
139




Fig. 20.3 Trajectory of total phosphorus (TP) versus chlorophyll-a (Chl-a) in Alte Donau during eutrophication and restoration (1987–2011). Years with macrophyte or algal domination are indicated by white and grey background respectively separated by a line equivalent to $\text{Chl-a} = 0.5\text{TP}$. TP and Chl-a as annual averages. Trophic delineations are indicated by dashed lines

resilience after perturbation. The starting point for intensive regrowth of macrophyte was given by an intervention into the hydrological regime of Alte Donau (Fig. 20.2). From 2002 onwards every spring the water level was lowered in the range of 15–30 cm to improve the light availability at the lake bottom at the beginning of the growing season and to imitate partly the former natural hydrological dynamic. The watermilfoil started to grow as well from planted areas as spontaneously in other lake areas. Other species that have been planted showed initially good growth but were more and more suppressed by shading due to the dense stands of *Myriophyllum spicatum*.

Seven years after the Riplox treatment the system switched back to the macrophyte dominated clear-water state. The response of macrophyte biomass to the increase and subsequent decrease in TP concentrations is shown in Fig. 20.4. The trajectory indicates large hysteresis in the loss and re-colonization of under-water vegetation. Management plans to stabilise rehabilitated ecosystems need to take hysteretic behaviour and return time into consideration.

Water quality in terms of total phosphorus and chlorophyll-a is now, 22 years after the forward shift, better than it was back in 1987 (Fig. 20.3, details in Chap. 6), but the quantity and composition of submerged vegetation is still quite different.




Fig. 20.4 Hysteresis in the decline and recovery trajectories of macrophyte biomass versus total phosphorus concentration (TP) in the open water during the eutrophication and rehabilitation phases between 1987 and 2011. Macrophyte biomass as in Fig. 20.2, TP as in Fig. 20.4

Macrophyte beds are dominated up to more than 90% by *Myriophyllum spicatum*, whereas the ground-covering charophytes are under-represented.

The relative contribution of different plant groups to total macrophyte biomass is documented for the period before eutrophication with macrophyte domination for the year 1987 and for the years 1993, 2002 and 2015 following treatment (Fig. 20.5, see also Fig. 20.2).

Myriophyllum spicatum is characterised by a very successful growth strategy but the dominance was also triggered by additional effects. At the beginning of the 1990th the plant stands regularly died back in the winter months. By the end of the 1990th the *Myriophyllum* stands started to overwinter as green plants, possibly due to a series of mild winters in the last decade that might be an effect of global warming. The impact of global change on Alte Donau was verified by the correspondence between the climate signal (North Atlantic Oscillation Index) and water temperature (WT) in winter and early spring, the increase of 1.52 °C per decade for surface water temperature in April and the prolongation of warm period in summer (Chap. 11).

Two years after full recovery of macrophytes and due to intensive photosynthesis of macrophytes and also of phytoplankton (algal primary production see Chap. 10) and low ground water influx especially into the main basin of Obere Alte Donau, pH-values tended to rise up to 10 in the summer months. Water from the impoundment




Fig. 20.5 Relative shares of different plant groups in the years 1987 (last documented good status before the eutrophication phase), 1993 (nearly total loss of macrophyte vegetation), 2002 (7 years after the Riplox treatment at the starting point of the periodical water level lowering in spring) and now (2015)

Neue Donau with a higher buffer capacity was used to exchange with water from Alte Donau to overcome this problem. This measure was done regularly from 2006 up to now in summer or autumn, exchanging quantities of 1.5–4.5 Mio m³ within several weeks. The buffer capacity in Alte Donau could thus be raised sufficiently and the pH-values settled between 8 and 9.

A special soil filter, which is in use since 2016, was constructed in the northwestern part of Alte Donau (Wasserpark) to find a sustainable solution to stock up the main basin of Obere Alte Donau with calcium and to raise alkalinity. This soil filter can be fed with water from Neue Donau. Phosphorus, suspended materials and chlorophyll-a are reduced when passing the soil filter. The water from Neue Donau is marked by higher calcium concentrations and by higher alkalinity than the water of Alte Donau. But the special feature of the soil filter is, that the Ca content can be stocked additionally by passing a reservoir that is filled with calcium carbonate before entering the system of Alte Donau. The filter allows a constant discharge of 2500 m³ per day, which should be sufficient to raise buffer capacity of Alte Donau permanently. This prototype of a soil filter was planned and built within an EU-Life project (EU-Life 12 ENV/AT/000128).

Because of the overall dominance of the high-growing species *Myriophyllum spicatum* intensive mowing is necessary to ensure bathing and other activities (Chaps. 8 and 19). Mowing the macrophytes also enhances the availability of light in deeper zones stimulating the growth of the low-growing vegetation. To support the proliferation of stoneworts intensive plantings have been done (Chap. 8). In contrast to helophyte planting, which turned out to be feasible with reasonable effort (Chap. 18), planting and re-establishing dense stonewort stands was a great



Fig. 20.6 Macrophyte cover in the Alte Donau in 1987 after Löffler (1988) – light grey: charophytes, dark grey: *Myriophyllum* spp. and *Potamogeton* spp. – serving as objective of the macrophyte management

challenge. As well-documented by an automatic underwater video-trap (Chap. 8) fish effectively hindered the growth of charophytes by grazing (amur and rudd) or by digging out the new planted stoneworts (bream and carp, fish assemblages in Chap. 15). In order to allow a successful re-establishment of charophytes in the Alte Donau fish management has to be optimised. Charophytes can preserve a good water quality equally well as the currently dominating high-growing vegetation (Van den Berg et al. 1998), therefore the stoneworts are an ideal aquatic weed group for water quality management of intensively used urban lakes. The conditions in the Alte Donau in the 1980ies (Löffler et al. 1988) can serve as a benchmark for a sufficient cover of charophytes in a mesotrophic urban lake (Fig. 20.6).




Fig. 20.7 Dependence of total fish catch (tons) on phytoplankton chlorophyll-a (mg m⁻³) for the post-restoration years (1998ff.). Chl-a data from DWS-Hydro-Ecology, total fish catch data from Austrian Fishery Association

212 In the time frame of the restoration process of Alte Donau a lot of efforts have
213 been made to trigger changes in the fish community. Structure of fish assemblages
214 was modified by biomanipulation in 1998 in 'Kaiserwasser' (Chap. 15) and changes
215 in fisheries management and practice. Predators like pike-perch, asp and later on
216 pike were intensively stocked to reduce the planctivorous and non-predatory fish
217 population (Chap. 15). Since macrophyte stands recovered pike is the main predator
218 in Alte Donau. Restructuring of the fish community promoted also changes in the
219 zooplankton assemblage (Chap. 11), stimulating the upgrowth of mainly copepods
220 and some large species of macrophyte-habitat associated cladocerans.

221 As a result of a consequently performed fish management, total fish catches have
222 declined considerably and became significantly dependent on chlorophyll-a con-
223 centrations ($r^2 = 0.81$, $p < 0.001$, $n = 15$, see also Chap. 15) since 1998, the post-
224 restoration period (Fig. 20.7.). A recent study of the fish biocenosis according to the
225 European Water Framework Directive methodology (Gassner et al. 2013) attested
226 the Alte Donau a "good ecological status" (Gassner et al. 2014). Since 2007 phyto-
227 plankton is also assessed according to the WFRD and the results also indicate the
228 "good ecological status".

20.6 Conclusions

229

It can be concluded that suitable concepts and methods exist to establish and stabilise a macrophyte-dominated state in shallow lakes. Successful re-introduction of the aquatic macrophyte vegetation must be accompanied by adjusting the overall nutrient balance and the protection of the young aquatic plants against the negative influence of fish (herbivory, enhanced turbidity, nutrients) and waterfowl (herbivory, nutrients). Public awareness for the importance of 'aquatic weeds' has to be raised because of the potential of macrophytes to compete successfully with phytoplankton. Using macrophytes to keep water bodies transparent may include macrophyte management and short term restrictions to water sports.

According to Heintz (2004) and Taylor and Goldstein (2010) the management of water resources requires a new paradigm, the concept of sustainability. Restoration of the eutrophied urban lake Alte Donau was achieved by the implementation of an integrated management plan. Seven years after chemical treatment and after introduction of biomanipulation and other ecotechnical measures a switch back to a macrophyte dominated clear water stable state was observed. Routine monitoring of water quality and hydrology was essential for fine tuning and performing of a continuous management. Macrophyte domination and associated with this – a good water quality – could be stabilised for more than 20 years, demonstrating the quality of our management. In case of the urban lake Alte Donau the concept of sustainability is suitable and successful. Actually, in the frame work of the EU-Life project – Urban Lake Alte Donau (Life 12 ENV/AT/000128), two plans are compiled for maintaining and ensuring the "good ecological status". The Integrated Lake Management Plan will include all management measures and a plan for monitoring. The Risk Management Plan will give advices how to reduce the vulnerability of the ecosystem to effects like climate change and other anthropogenic pressures. This plan will be made accessible to authorities in other cities and can serve as a model for management of urban lakes in Europe.

References

257

- Bailey P, Boon P, Morris K (2002) Managing nutrients in floodplain wetlands and shallow lakes. River management technical update No. 2, Land & Water Australia, Canberra, 8pp
- Berg S, Jeppesen E, Sondergaard M (1997) Pike (*Esox lucius* L.) stocking as a biomanipulation tool I. Effects on the fish population in Lake Lyng, Denmark. *Hydrobiologia* 342–343:311–318
- Berger J, Schagerl M (2003) Allelophytic activity of *Chara aspera*. *Hydrobiologia* 501:109–115
- Birch S, McCaskie J (1999) Shallow urban lakes: a challenge for lake management. *Hydrobiologia* 95/396:365–377
- Björk S (1978) Restoration of degraded lake ecosystems. *Reports of the Institute of Limnology*. University of Lund, 24pp


- Björk S (1994) Restoration of lakes through sediment removal – Lake Trummen, Sweden. In: Eiseltova M (ed) Restoration of lake ecosystems. A holistic approach, vol 32. IWRB Publications, pp 130–140
- Bowmer KH, Mitchell DS, Short DL (1984) Biology of *Elodea canadensis* MICHX. and its management in Australian irrigation systems. *Aquat Bot* 18:231–238
- Cappiella K, Schueler T (2001) Crafting a lake protection ordinance. *Water Prot Tech* 3:751–768
- Carvalho L (1994) Top-down control of phytoplankton in a shallow hypertrophic lake: little Mere (England). *Hydrobiologia* 275–276:53–63
- Crisman TL, Mitraki C, Zalidis G (2005) Integrating vertical and horizontal approaches for management of shallow lakes and wetlands. *Ecol Eng* 24:379–389
- Dokulil MT (2014a) Environmental impacts of tourism on lakes. In: Ansari AA, Gill SS (eds) Eutrophication: causes, consequences and control, vol 2. Springer, pp 81–88, ISBN 978-94-007-7813-9
- Dokulil MT (2014b) Old wine in new skins: eutrophication reloaded: global perspectives of potential amplification by climate warming altered hydrological cycle and human interference. In: Lambert A, Roux C (eds) Eutrophication: causes, economic implications and future challenges. Nova Publisher, Hauppauge, ISBN 978-1-62808-498-6
- Gassner H, Achleitner D, Bruschek G, Mayrhofer K, Frey I (2013) Leitfaden zur Erhebung der Biologischen Qualitätselemente – Teil B1 – Fische. BMFLUW, Vienna (ed.) <http://wisa.lebensministerium.at>
- Gassner H, Luger M, Dachleitner D, Pamminer-Lahnsteiner B (2014) Alte Donau (2013) – Standardisierte Fischbestandserhebung und Bewertung des fischökologischen Zustandes gemäß EU-WRRL. Bericht, 34 Seiten. Bundesamt für Wasserwirtschaft, Institut für Gewässerökologie, Fischereibiologie und Seenkunde, Scharfling, <http://www.baw.at/index.php/igf-publikationen/170-igf-publikationen-nach-jahr.html>
- Gopal B, Goel U (1993) Competition and allelopathy in aquatic plant communities. *Bot Rev* 59:155–210
- Gross E, Hilt S (2005) Probleme mit submersen Makrophyten in Flachseen Deutschlands. http://www.f.igb-berlin.de/workshop_2005/Gross_Hilt.pdf
- Hansel-Welch N, Butler MG, Carlson TJ, Hanson MA (2003) Changes in macrophyte community structure in Lake Christina (Minnesota), a large shallow lake, following biomanipulation. *Aquat Bot* 75:323–337
- Hansson MA, Butler MG (1994) Responses of plankton, turbidity, and macrophytes to biomanipulation in a shallow prairie lake. *Hydrobiologia* 200–201:317–327
- Heintz HT (2004) Applying the concept of sustainability to water resources management. *Water Resour Update* 127:6–10
- Hilt S (2005) Schaffung von Bedingungen zur natürlichen Wiederansiedlung von submersen Makrophyten in Flachseen. http://www.f.igb-berlin.de/workshop_2005/Hilt.pdf
- IGB – Leibnitz-Institut für Gewässerökologie und Binnenfischerei (2005) Workshop des DGL<AK “Flachseen”, limnologische Station der TU München. http://www.f.igb-berlin.de/workshop_2005
- Irfanullah HM, Moss B (2004) Factors influencing the return of submerged plants to a clear-water, shallow temperate lake. *Aquat Bot* 80:177–191
- Janauer GA (1995) Makrophytenmanagement in Donaualtarmen – Wissenschaftliche Zustandsanalyse und Managementkonzept (Macrophyte management in Danube oxbows. – Scientific status assessment and management concept). Technical Report, Verbundgesellschaft, Wien, 111pp
- Janauer GA, Pall K (1998) Donaualtarme als Lebensraum. Schriftenreihe der Forschung im Verbund 37, Österreichische Elektrizitätswirtschafts-Aktiengesellschaft, Wien, 46pp
- Janse JH, van Donk E, Aldenberg T (1998) A model study on the stability of the macrophyte-dominated state as affected by biological factors. *Water Res* 32:2696–2706
- Körner S (2002) Submerse Makrophyten – wichtig für die Seentherapie in Deutschland? *Wasser & Boden* 9:38–41

- Lau SSS, Lane SN (2002) Nutrient and grazing factors in relation to phytoplankton level in a eutrophic shallow lake: the effect of low macrophyte abundance. *Water Res* 36:3593–3601
- Madsen JD, Adams MS, Ruffier P (1988) Harvest as a control for Sago Pondweed (*Potamogeton pectinatus* L.) in Badfish Creek, Wisconsin: frequency, efficiency and its impact on the stream community oxygen metabolism. *J Aquat Plant Manag* 26:20–25
- Morscheid H, Maehlmann J (2005) Künstliche (Wieder-)Ansiedlung von Makrophyten. http://www.f.igb-erlin.de/workshop_2005/Morscheid_Maehlmann.pdf
- NALMS – North American Lake Management Society (1988) Lake and reservoir restoration guidance manual EPA. 440/5-88-002
- Naselli-Flores L (2008) Urban Lakes: ecosystems at risk, worthy of the best care. In: Sengupta M, Dalwani R (eds) *Proceedings of Taal2007: the 12th World Lake conference*, pp 1333–1337
- Perrow MR, Meijer M-L, Dawidowicz P, Coops H (1997) Biomanipulation in shallow lakes: state of the art. *Hydrobiologia* 342–343:355–365
- Pieterse AH and Murphy KJ (1990) *Aquatic weeds. The ecology and management of nuisance aquatic vegetation*. Oxford University Press, Oxford, 593 pp
- Riis T, Hawes I (2002) Relationship between water level fluctuations and vegetation diversity in shallow water of New Zealand lakes. *Aquat Bot* 74:133–148
- Schueler T, Simpson J (2001) Why urban lakes are different. *Water Prot Tech* 3(4):747–750
- Singh KP (1982) Lake restoration methods and feasibility of water quality management. State Water Survey Division, Surface water section, University of Illinois, *Report 301*, 53pp
- Sorensen HA (1996) Managing urban lakes: an integrating experience. *Water Resour Dev* 12:437–446
- Taylor T, Goldstein R (2010) Sustainable Water research management 3: case studies on new water paradigm. Report 1020587, Electronic Power Research Institute, Palo Alto, 172pp
- Traut AH, Hostetler ME (2004) Urban lakes and waterbirds: effects of shoreline development on avian distribution. *Landsc Urban Plan* 69:69–85
- Van den Berg MS, Coops H, Meijer M-L, Scheffer M, Simons J (1998) Clear water associated with a dense *Chara* vegetation in the shallow and turbid Lake Veluvemeer, The Netherlands. In: Jeppesen E, Sondergaard M, Sondergaard M, Christoffersen K (eds) *The structuring role of submerged macrophytes in lakes*. Springer, New York, pp 339–352
- Van der Does J and Frinking LJ (1993) Restoration of Lake Geerplas by reduction of external phosphorus loading and dredging of phosphorus enriched upper sediment layer (Abstract) *Verhandlungen Internationale Vereinigung Limnologie* 25, 614
- Welch EB, Schriever GD (1994) Alum treatment effectiveness and longevity in shallow lakes. *Hydrobiologia* 275:423–431
- Wium-Andersen S (1987) Allelopathy among aquatic plants. *Arch Hydrobiol* 27:167–172
- Wychera U, Humpesch UH (2002) Translocation and growth experiments with the macrophyte *Potamogeton lucens* L. in Traunsee (Austria), with respect to industrial tailings. *Water Air Soil Pollut Focus* 2:109–116
- Xu FL, Tao S, Xu ZR (1999) The restoration of riparian wetlands and macrophytes in Lake Chao, an eutrophic Chinese lake: possibilities and effects. *Hydrobiologia* 405:169–178

Author Queries

Chapter No.: 20 0003946999

Queries	Details Required	Author's Response
AU1	Please confirm the corresponding author.	
AU2	Please confirm the "Sentence "Methods have been summarised...." for correctness".	
AU3	References Löffler et al. (1988), Löffler (1988) are cited in the text, but not in the list. Provide a citation for them or delete these citations from the text.	
AU4	Please provide publisher location for the references Björk (1994) and Dokulil (2014a).	



1st ed. 2018, VIII, 310 p. 181 illus., 54 illus. in color.

Printed book

Hardcover

139,99 € | £119.99 | \$169.99

^[1]149,79 € (D) | 153,99 € (A) | CHF 165,50

eBook

118,99 € | £95.50 | \$129.00

^[2]118,99 € (D) | 118,99 € (A) | CHF 132,00

Available from your library or
[springer.com/shop](https://www.springer.com/shop)

MyCopy ^[3]

Printed eBook for just

€ | \$ 24.99

[springer.com/mycopy](https://www.springer.com/mycopy)

Martin T. Dokulil, Karl Donabaum, Katrin Teubner (Eds.)

The Alte Donau: Successful Restoration and Sustainable Management

An Ecosystem Case Study of a Shallow Urban Lake

Series: Aquatic Ecology Series

- Provides a variety of strategies to restore and manage deteriorated recreational urban waters.
- Broadens the understanding of the importance of all trophic levels of the food web for successful rehabilitation and implementation of landscape architecture for an attractive recreational site
- Clearly demonstrates that long-term efforts are essential for sustainable recovery

Here we report on a 25-year long-term sequence of measures to return a deteriorated recreational urban lake, Alte Donau in Vienna to acceptable water quality. Metropolitan waters require focused ecosystem management plans and intensive in-lake efforts. We explored physico-chemical conditions, food web from viruses to fish and water birds, the sediments, the littoral zone and the catchment, management and urban planning, and global warming. Several restoration techniques were tested and critically evaluated. The final management plan was based on bi-stable theory. During the recovery phase, numerous surplus adjustments had to be implemented to secure sustainable achievement.

Lifelong 40% discount for authors



Order online at [springer.com](https://www.springer.com) / or for the Americas call (toll free) 1-800-SPRINGER / or email us at: customerservice@springernature.com. / For outside the Americas call +49 (0) 6221-345-4301 / or email us at: customerservice@springernature.com.

The first € price and the £ and \$ price are net prices, subject to local VAT. Prices indicated with [1] include VAT for books; the €(D) includes 7% for Germany, the €(A) includes 10% for Austria. Prices indicated with [2] include VAT for electronic products; 19% for Germany, 20% for Austria. All prices exclusive of carriage charges. Prices and other details are subject to change without notice. All errors and omissions excepted. [3] No discount for MyCopy.

Part of **SPRINGER NATURE**